



Evaluation of Survival Rate in Patients with Diabetic Foot Ulcer with Comorbid Conditions in Tertiary Care Hospital

Sk. Salma Sultana^{1*}, Sk. Musfira Kauser², S. Sirisha Reddy³, Md. Muddasseer⁴, K. Venkata Krishna⁵, G.V. Prakash⁶, Kudipudi Harinadha Baba⁷

¹Department of Pharmacology, Narayana Pharmacy College, Chinthareddypalem, Nellore-524002.

²Pharm D, Narayana Pharmacy College, Chinthareddypalem, Nellore-524002.

³Pharm D, Narayana Pharmacy College, Chinthareddypalem, Nellore-524002.

⁴Pharm D, Narayana Pharmacy College, Chinthareddypalem, Nellore-524002.

⁵Pharm D, Narayana Pharmacy College, Chinthareddypalem, Nellore-524002.

⁶Department of General Surgery, Narayana Medical College, Chinthareddypalem, Nellore -524002.

⁷Department of Pharmaceutical Analysis, Narayana Pharmacy College, Chinthareddypalem, Nellore - 524002.

ABSTRACT

Aim: Diabetic foot ulcer is the most common complication of diabetic foot ulcer and has a significant lifethreatening impact on the patients especially patients with other comorbidities predominantly diabetes mellitus. Therefore, the purpose of this study is to estimate the survival rate in various patients suffering from DFU associated with comorbidities such as hypertension, Diabetes Mellitus, Cardiovascular diseases and other complications. **Materials and methods:** An observational retrospective cohort study was conducted on approximately 120 patients in general surgery department of Narayana medical hospital over a period of 6 months October 2022 to May 2023. Ethical clearance also has been acquired from the Institutional review committee at Narayana Medical College. **Results:** Among the 120 patients observed during the study, DFU along with only DM comorbidity were recorded in highest number i.e., 31 (35.83%). 16.66% and 11.66% DFU patients were suffering from DM along with hypertensions and CVS diseases respectively. Maximum no.of deaths were recorded in DFU patients with CVS diseases (43.75%) when compared to other comorbid conditions. Based on the data collected, 86.6% survival risk probability was observed in patients with CVS diseases indicating lower survival chances. **Conclusion:** Based on the available data collected during the study, after assessing the data and acquiring the results, it was concluded that patients with DFU associated with CVS diseases comorbidities has lower survival based on over survival rate.

Keywords: Diabetic foot ulcer, Survival Rate, Retrospective cohort study, comorbid conditions

ARTICLE INFO

Corresponding Author

Sk. Salma Sultana
Department of Pharmacology,
Narayana Pharmacy College,
Chinthareddypalem, Nellore-524002..

Article History

Received : 12 Mar 2023
Revised : 14 May 2023
Accepted : 19 June 2023
Published : 24 Aug 2023

Copyright© 2023 The Contribution will be made Open Access under the terms of the Creative Commons Attribution-NonCommercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0>) which permits use, distribution and reproduction in any medium, provided that the Contribution is properly cited and is not used for commercial purposes.

Citation: Sk. Salma Sultana, et al. Evaluation of Survival Rate in Patients with Diabetic Foot Ulcer with Comorbid Conditions in Tertiary Care Hospital. A. J. Med. Pharm, Sci., 2023; 11(1): 24-28.

CONTENTS

1. Introduction.....	24
2. Methodology.....	25
3. Results and Discussion.....	25
4. Conclusion.....	28
5. References.....	28

1. Introduction

Diabetic foot ulcer is the most common complication in patients with DM^[1]. DFU is caused by bacterial invasion which results in infection and decay, these ulcers occur in any part of the body particularly in the distal part of the lower leg^[2]. There are three types of DFU namely, neuropathic

ulcers (diabetic ulcers), arterial leg ulcers (ischemic ulcers) and venous leg ulcers (stasis ulcers)^[3]. Depending upon the severity, based on Wagner's classification, DFU is classified into Grades. No ulcer, superficial ulcer, Deep ulcer, ulcer with a bcess, localized gangrene and extensive

gangrene are classified as grade 0,1,2,3,4 and 5 respectively^[4].

Epidemiology: During their lifetime 15 % of them are suffered from DFU.[17] In DFU patients 85% of the amputation takes place. The proportion of DFU includes that 54%, 34%, and 10% of the individuals are suffered from non-ischemic, ischemic and purely ischemic ulcers^[5].

Etiology

DFU is mainly caused by bacteria (both gram positive and gram negative) predominantly *S. aureus*, *E. coli* and *Klebsiella* species^[6]. Other causative factors include peripheral neuropathy, high foot plantar pressure and trauma. In peripheral neuropathy, loss of pain sensation may lead to unnoticed repetitive injuries at pressure points facilitating the formation of ulcer^[7]. Some of the contributive factors include Atherosclerosis and diabetes^[7].

Pathophysiology

In neuropathic condition, predominantly sensory neuropathy, the unnoticed trauma may progress into lesions due to loss of pain sensation. In the lesion, tissue necrosis and callus formation develop into the cavities filled with serous fluid that leads into the formation of ulcers. Skin infections, abscess formation, sepsis and gangrene are some of the complications of DFU^[8]. In vasculopathy (microangiopathy and macroangiopathy), arterial occlusion occurs either due to increased basement membrane thickness or atherosclerosis of arteries that leads to decreased blood circulation to the ulcer^[8].

Symptoms

Common symptoms of DFU include skin discoloration (darkened skin), pain, numbness, loss of hair. Instead, ischemic ulcer do not have callus formations but are represented with gangrene at regions such as ankles and toes^[9]. Stasis ulcer has presented with hardened skin and well in around the ulcer site. decreased circulation to the injury site which in turn prolongs healing of an ulcer^[10].

Diagnosis

They are diagnosed as foot ulcer by examining the local site of the ulcer including parameters pertaining to size, region, type of discharge and severity of the ulcer^[11]. They are diagnosed using X rays, MRI scan, CT scan, blood test, high foot plantar examination etc.,^[12]

Treatment^[13]

Based on severity of the ulcer, treatment using medications include, Surgical methods include Incision, Debridement, Wound lavage, wound closure and Amputation. Debridement is carried out by the removal of devitalized tissue. It is the most effective method as it reduces the risk and shortens the healing time^[14]. Primary closure is the closure of the wound at the end of the surgery prolonging healing time for infection^[14]. Amputation is removing the infected portion of the limb or whole limb to save the rest of the portion^[14].

Table 1

Severity	Treatment
Mild to moderate	Cephalexin 500 mg-QID, Amoxicillin/clavulanate –875/125mg
Moderate severe	Clindamycin 300 mg TID, Ampicillin/sulbactam 3g TID

Life threatening	Imipenem/ cilastin 500 mg QID, clindamycin+ Ampicillin+ Tobramycin (900mg+5.1mg+500mg) QID
------------------	--

2. Methodology

Inpatient department [IPD] of general surgery in tertiary care hospital i.e. Narayana medical college and hospital in Nellore region.

Study design

It is an Observational Retrospective cohort study.

Study population

In this study the patients who had diagnosed with diabetic foot ulcer and other comorbid conditions of about 120 patients were included for follow up study period October 2022-May 2023.

Study criteria

Inclusion criteria

Patient with DFU and other comorbid conditions
Aged ≥ 18 years old
Continuous existence of diabetic foot lesion for a minimum of 4 weeks

Exclusion criteria

Severe Anemia which is not caused by an infection

Pregnancy

Unavailable medical records and unequipped data
Simultaneous participation in other investigational trials/previous participation in trials

Study material:

Data were collected by interview in the patients and from the data collection proforma, Case sheets of the patients.

Study Variables

The study variables include patient had any other chronic problems, Family history and socio demographic details [Age, Sex, Place of birth, Personal history, Habits and Employment status] and comorbid condition Includes DM, CVD, HTN, Renal diseases, Respiratory diseases, Thyroid conditions and others. The main outcome of the study was to determine the survival rate of Diabetic foot ulcer and other comorbid conditions. The risk factors for development of DFU includes in this study were Age, Sex, Diabetes duration, any foot appearance {Foot deformities, Peripheral Neuropathy, Peripheral artery diseases [PAD]}, HTN, Fasting plasma glucose, HbA1C, LDL.

Statistical Analysis

All data was collected and analysis was performed. It includes descriptive analysis where in patients clinical condition and various variables were analyzed for their respective frequencies and percentages. Statistical analysis was used to determine whether there is a significant difference between the survival rate of patients with DFU in different comorbid conditions. The statistical analysis used is paired t-test that is to determine the significant difference between the no. of patients affected and no. of patients survived.
 $t = \frac{d}{s/\sqrt{n}}$

3. Results and Discussion

- In this study a total of 120 patients with DFU along with comorbid conditions who visited the inpatient department of general surgery at tertiary care hospital were included.

- The age groups of the subjects were taken from 20-90 years among them the age group between 41-50 years [30] are more affected with DFU the percentage was found to be 25%, 71-80 years age group subjects i.e., 29 of them are affected with the DFU [24.1666%], 23 subjects [19.1666%] were affected with DFU between the age group of 61-70, 21 subjects [17.5%] between the 51-60 years, 11 subjects [9.1666%] between 31-40 years, 3 [2.5%] of them between 21-30 and 81-90 age groups [Table-1]
- Subject characteristics based on ulcer size were shown in Table- 2. In this study majority of subjects i.e 86 [71.6%] had the ulcer size within the range of 0-10 cms, ulcer size between the range of 11-20cms, ulcer size between the range of 50-110cms were found in least subjects.
- According to the study, Patients with DFU along with different comorbid condition has shown in Table-3. 3 [25.83%] subjects had only DM, investigational methods used to detect other comorbid conditions has revealed that 20 [16.66%] had HTN along with DM, 14 [11.66%] had cardiovascular diseases [CAD, Atherosclerosis], 6 [5%] had renal [CKD, AKI] and respiratory diseases [Asthma, COPD], 3 [2.5%], 2 [1.66%], 38 [31.66%] had psychiatry, thyroid, other disease conditions [SLE, LEA] respectively.
- Among 120 patients most of them 37 [30.83%] had undergone surgery of debridement to prevent the severity of the wound of the ulcer, Ray amputation, Below knee amputation, Toe amputation has been performed in about 26 [21.6%], 14 [11.6%], 96 [6.66%] patients respectively to remove the infected toe or limb so, that the further, posterior spread of the wound would be ceased, Fasciotomy has been performed in about 8-12 patients [7.5%] to relieve wound pressure, Primary suturing, Tendelberg procedure, Disarticulation has been performed as supportive treatment along with conservative management. [Table-4]
- Among the admitted patients with DFU, majority of the subjects 7 [43.25%] with cardiovascular diseases recorded as death, 4 deaths [25%] were recorded in subjects with HTN along with DM, 2 death [12.5%] were recorded in subjects with thyroid diseases and subjects with DM only. [Table-5]

- The Survival probability of patients with CVD was found to be 0.5 [50%], DM probability was 0.935 [93.5%], DM with HTN was 0.8 [80%], Respiratory and thyroid conditions had the survival probability of 0 [100%], Renal, Psychiatry, other disease conditions had the survival probability of 0.834 [83.4%], 0.667 [66.7%], 0.9737 [97.37%] respectively. [Table-6]







Table 2

No. of patients	Only DM	DM with HTN	DM with Respiratory diseases	DM with CVS diseases	DM with Renal diseases	DM with Psychiatry diseases	DM with Thyroid diseases	DM with other diseases
Effected	31	20	6	14	6	2	3	38
Survived	29	16	6	7	5	2	2	37
Died	2	4	0	7	1	0	1	1

From the above data, the t-value calculated is 2.3665, Whereas t-tabled value at 5% LOS with 7 DOF is 1.895.

Since $t_{cal} > t_{tab}$, We reject H_0 and We accept H_1 Where, H_0 – There is no significant difference between the no. of patients affected & no. of patients survived

Table-3: Subjects According to their Age Groups

AGEGROUP	NO.OFSUBJECTS	PERCENTAGE
21-30	3	2.5
31-40	11	9.1666
41-50	30	25
51-60	21	17.5
61-70	23	19.1666
71-80	29	24.1666
81-90	3	2.5%

Table-4: Classification of Subjects According to their Ulcer Size

ULCERSIZE	NO.OFSUBJECTS	PERCENTAGE
0-10CM	86	71.6
11-20CM	16	13.333
21-30CM	6	5
31-40CM	3	2.5
41-50CM	1	0.8333
51-60CM	2	1.666
61-70CM	1	0.8333
71-80CM	2	1.666
81-90CM	0	0
91-100CM	1	0.8333
101-110CM	2	1.666

Table-5: Shows Data of Subjects with DFU in Different Comorbid Conditions



COMORBIDCONDITION	NO.OFSUBJECTS	PERCENTAGE
OnlyDM	31	25.83
DMalongwithHTN	20	16.66
DMalongwithrespiratorydisease	6	5
DMalongwithCVSdisease	14	11.66
DMalongwith renaldisease	6	5
DMalongwith psychiatrydisease	3	2.5
DMalongwiththyroidcondition	2	1.66
DMalongwith other conditions	38	31.66

Table-6:Shows Deaths in Subjects Suffering From DFU

Disease	No.ofsubjects	No.ofsubjectsdeath	Percentage
OnlyDM	31	2	12.5
DMwith HTN	20	4	25
DMwithrespiratorydisease	6	0	0
DMwithCVSdisease	14	7	43.75
DMwithrenaldisease	6	1	6.25
DMwithpsychiatry	3	0	0
DMwiththyroidcondition	2	1	6.25
DMwithotherconditions	38	1	6.25

Table-7: Data Shows Survival Probability of Individual Comorbid Condition

Comorbidcondition	No.ofsubjectsaffected	No.ofsubjects survived	Probability
DM	31	29	0.9354[93.54%]
DMwith HTN	20	16	0.80[80%]
Cardiovascular diseases	14	7	0.50[50%]
Respiratory diseases	6	6	0[100%]
Thyroid diseases	2	2	0[100%]
Renal diseases	6	5	0.834[83.4%]
Psychiatry diseases	3	2	0.667[66.7%]
Other conditions	38	37	0.9737[97.37%]



4. Conclusion

Two years overall survival probability of DFU was 0.866[86.6%]. Patients with cardiovascular diseases are high risk for development of foot ulcer have lower survival probability when compared with the other comorbid conditions [Renal diseases, DM with HTN, Thyroid condition, DM, Respiratory, Psychiatry diseases].

5. References

- [1] Oliver TI, Mutluoglu M. Diabetic foot ulcer. Rosyid FN. Etiology, pathophysiology, diagnosis and management of diabetics' foot ulcer. International Journal of Research in Medical Sciences. 2017 Oct, 5(10):4206-13.
- [2] AlexandrescuVa, Deleeuw P, KovandaJs. Ischemic and Venous Wound Identification: What We Look For. Crit Limb Ischemia. 2017;16(5):1-6.
- [3] Shah P, Inturi R, Anne D, Jadhav D, Viswambharan V, Khadilkar R, Dnyanmote A, Shahi S, Shah PP, Inturi Jr R, Dnyanmote AS. Wagner's classification as a tool for treating diabetic foot ulcers: Our observations at a suburban teaching hospital. Cureus. 2022 Jan 22; 14(1).
- [4] Chahine EB, Harris S, Williams R. Diabetic foot infections: an update on treatment. US Pharm. 2013;38(4):23-6.
- [5] Rosyid FN. Etiology, pathophysiology, diagnosis and management of diabetics' foot ulcer. International Journal of Research in Medical Sciences. 2017 Oct; 5(10): 4206-13.
- [6] Syafril S. Pathophysiology diabetic foot ulcer. In IOP Conference Series: Earth and Environmental Science 2018 Mar 1 (Vol. 125, No.

1, p. 012161). IOP Publishing. Dendale A. Derm Net New Zealand.

- [7] Ashcroft E. Recognizing and treating venous stasis ulcers. Deepes Tissues: Wounds Australia newsletter. 2017 Sep 1:36-9.
- [8] Chahine EB, Harris S, Williams R. Diabetic foot infections: an update on treatment. US Pharm. 2013;38(4):23-6.