



World Journal of Pharmacy and Biotechnology

ISSN: 2349-9087 | www.pharmaresearchlibrary.com/wjpb

W. J. Pharm. Biotech., 2015, 2(1): 55-61

DOI: https://doi.org/10.30904/j.wjpb.2015.2584



Laccase–Production and It’s Appliacion

Shruti S. Patel*, Sanjay N. Parekh

Shree Ram Krishna Institute of Computer Education & Applied Science, Surat, India

ABSTRACT

Laccase belongs to the small group of enzymes called the blue multi copper oxidases. Laccase is widely distributed in higher plants, fungi, bacteri. By fermentation both submerged and solid state fermentation laccase can be produced by using various carbon and nitrogen sources. Laccases have been subject of intensive research in the last decades due to their broad substrate specificity. In the recent years, their uses span from the textile to the pulp and paper industries, and food applications to bioremediation processes. More recently, they have found applications in other field such as in the design of biosensors. In this review, the occurrence, structure, production, and application of laccases will be discussed.

Keywords: Laccase, sources, Production, structure, applications, Pulp and paper industry, Food industry, Bioremediation, Biosensors

ARTICLE INFO

Corresponding Author

Shruti S. Patel
Shree Ram Krishna Institute of
Computer Education & Applied Science
Surat, India

Article History

Received : 18 Feb 2015
Revised : 28 Mar 2015
Accepted : 24 May 2015
Published : 29 June 2015

Copyright© 2015 The Contribution will be made Open Access under the terms of the Creative Commons Attribution-NonCommercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0>) which permits use, distribution and reproduction in any medium, provided that the Contribution is properly cited and is not used for commercial purposes.

Citation: Shruti S. Patel and Sanjay N. Parekh. Laccase – Production and it’s Appliacion. *W. J. Pharm. Biotech.*, 2015, 2(1): 55-61.

CONTENTS

1. Introduction.	55
2. Structure of Laccase.. . . .	55
3. Purification of Laccase.	56
4. Laccase Application.	57
5. Conclusion.	59
6. References	59

1. Introduction

Laccases (p-diphenol: dioxygen oxidoreductases; benzenediol dioxygen oxidoreductases; EC 1.10.3.2) are member of large blue copper proteins or the blue copper oxidases; other enzymes in this group are the plant ascorbate oxidase and the mammalian plasma protein ceruloplasmin [1,2]. Laccases are defined in the Enzyme Commission (EC) nomenclature as oxidoreductases which oxidize aromatic and nonaromatic compounds by a radical-catalyzed reaction mechanism and use molecular oxygen as an electron acceptor. Laccases catalyze the oxidation of a variety of phenolic compounds, as well as diamines and aromatic amines, with concomitant reduction of molecular oxygen to water [1]. The syringaldazine [4-hydroxy-3,5-

dimethoxy benzaldehyde azine] is considered the substrate oxidized only by laccase enzyme [1]. Thus, several organic compounds which contain hydroxyl, acid, or amino groups can act like substrates [3]. Recently a novel polyphenol oxidase with laccase like activity was mined from a metagenome expression library from bovine rumen microflora [3].

2. Structure of Laccase

Laccases often occur as isoenzymes or monomers that oligomerize to form multimeric complexes. Each isoenzyme has four copper atoms and is able to individually carry on the catalytic mechanism of laccases. The

Single-step laccase purification from *Neurospora crassa* takes place by using celite chromatography and 54 fold purification was obtained with specific activity of 333 U mg⁻¹ [17]. Laccase from LLP13 was first purified with column chromatography and then purified with gel filtration. Laccase from *T. versicolor* is purified by using ethanol precipitation, DEAE-Sepharose, Phenyl-Sepharose and Sephadex G-100 chromatography which is a single monomeric laccase with a specific activity of 91,443 U mg⁻¹ [41]. Laccase from *T. versicolor* is purified with Ion Exchange chromatography followed by gel filtration with specific activity of 101 U mL⁻¹ and 34.8-fold purification. Laccase from *Stereum ostrea* is purified with ammonium sulphate followed by Sephadex G-100 column chromatography with 70-fold purification. Laccase from fruiting bodies is purified with ammonium sulphate precipitation with 40–70% saturation and DEAE cellulose chromatography then 1.34 and 3.07 fold purification is obtained respectively [42].

4. Laccase Application

A few laccases are at present in the market for textile, food and other industries (Table 3), and more candidates are being actively developed for future commercialization [16]. Laccase is important because it oxidizes both the toxic and nontoxic substrates. Laccases have been used in a variety of industrial and environmental applications since they can oxidize a number of natural and synthetic substrates using oxygen, producing water as the only by-product. Laccases are versatile enzymes able to oxidize recalcitrant compounds like lignin, which makes them attractive for use in various biotechnological processes. A vast amount of industrial applications for laccases have been proposed. This enzyme is very specific, ecologically sustainable and a proficient catalyst. Applications of laccases are as follows.

Nanobiotechnology

Since laccases are able to catalyse electron transfer reactions without additional cofactors, their use has also been studied in biosensors or bioreporters to detect various phenolic compounds, oxygen or azides. Moreover, biosensors for detection of morphine and codeine, catecholamines, plant flavonoids, for determination of glucose, aromatic amines and phenolic compounds and also for electroimmunoassay have been developed. The Biotechnological application of such micropatterned surfaces: the production of islands of micrometer size of extracellular matrix, where the pattern of these islands could determine the position and distribution of bovine and endothelial cells [42]. Laccase covalently conjugated to a bio-binding molecule can be used as a reporter for immunochemical (ELISA, Western blotting), histochemical, cytochemical, or nucleic acid-detection assays.

Bioremediation and Biodegradation

Laccases have many possible applications in bioremediation. Laccase may be applied to degrade plastic waste having olefin units. Likely, an oxidation of the olefin units by the LMS, could initiate a radical chain reaction, leading to the disintegration of the plastic. Also this LMS can be used to degrade polyurethanes. LMS facilitated the

degradation of phenolic compounds (environmental hormones) from biphenol and alkylphenol derivatives and also the decomposition of fluorescent brighteners [17].

Laccase may also be used to eliminate odor emitted from places such as garbage disposal sites, livestock farms, or pulp mills. Also, they could be used for decolorizing dye house effluents that are hardly decolorized by conventional sewage treatment plants. In addition to dye house effluents, laccases can decolorize waste waters from olive oil mills and pulp mills by removing colored phenolic compounds. Another potential environmental application for laccases is the bioremediation of contaminated soils, as laccases and LMS are able to oxidize toxic organic pollutants, such as various xenobiotics, polycyclic aromatic hydrocarbons (PAHs), chlorophenols, and other contaminants.

The catalytic properties of laccases can be used to degrade such compounds. Phenolic compounds are present in wastes from several industrial processes, as coal conversion, petroleum refining, production of organic chemicals and olive oil production among others. Immobilized laccase was found to be useful to remove phenolic and chlorinated phenolic pollutants. Laccase was found to be responsible for the transformation of 2,4,6-trichlorophenol to 2,6-dichloro-1,4-hydroquinol and 2,6-dichloro-1,4-benzoquinone. LMSs have been also used to oxidize alkenes, carbazole, N-ethylcarbazole, fluorene, and dibenzothiophene. Isoxaflutole is an herbicide activated in soils and plants to its diketonitrile derivative, the active form of the herbicide: laccases are able to convert the diketonitrile into the acid. Laccase can be also used to reduce the concentration of synthetic heterocyclic compound such as halogenated organic pesticides in the soil. LMS has been extensively studied in the oxidation of recalcitrant PAHs, main components of several ship spills. In this sense, LMS is being included in several enzymatic bioremediation programs. *Cerrena unicolor* produces laccase in the low nitrogen medium which has the capability of reducing lignin content from sugarcane bagasse up to 36% within 24 h at 30°C.

Future Trends and Perspectives

This review summarizes the available recent important information about the properties, production of laccases and possible industrial and biotechnological use. Laccases are promising enzymes to replace the conventional chemical processes of several industries such as the pulp and paper, textile, pharmaceutical, and nanobiotechnology. However, one of the limitations to the large-scale application of laccases is the lack of capacity to produce large volumes of highly active enzyme at an affordable cost. The use of inexpensive sources for laccase production is being explored in recent times. Thus, efforts have to be made in order to achieve cheap over production of laccase from hosts, and also their modification by chemical means or protein engineering, to obtain more robust, active and less expensive enzymes. Another additional problem is the cost and toxicity of redox mediators. Further investigations should consider different and less polluting mediators such as the natural mediators produced by laccase in a bio-environment during lignin

degradation. The current development in laccase catalysis research and the design of mediators along with the research on its heterologous expression opens a wide spectrum of possible applications in the near future. Moreover, laccase can also offer a simple and convenient alternative to using peroxidases with H₂O₂, because laccases are available on an economically feasible scale. On the other hand, the development of an effective system for laccase immobilisation also deserves great attention. Immobilization could be achieved by chemical modification of the substrates. Hence, micropatterning, SAMs and LbL

techniques can be used to functionalise flat and curved surfaces in order to have specific adsorption. Laccase encapsulation with polyelectrolytes will be used as a microreactor for catalytic reactions by changing the permeability properties of the capsule wall. Since the general goal is to obtain stable catalysts with long life times and low cost, the combination of these techniques will enhance:

- a. The adsorption of laccase on a suitable substrate,
- b. The lifetime of the laccase activity and
- c. Reutilisation of the substrate/laccase product.

Table 1: Sources of Laccase Enzyme And its Physiological role

	sources	Physiological role	References
Plant	<i>Rhus vernicifera</i> , cabbages, turnips, beets, apples, asparagus, potatoes, pears, sycamore maple poplar tobacco , peach loblolly pine	<ul style="list-style-type: none"> • oxidize monolignols in the early stages of lignifications • involved in the first steps of healing in wounded leaves • in defense mechanism against external conditions 	[6,1,7,8,9,10,11,12]
Fungi	<i>Monocillium indicum</i> <i>Phanerochaete chrysosporium</i> , <i>Theiophora terrestris</i> , <i>Lenzites, betulina</i> , <i>Phlebia radiate</i> , <i>Pleurotus ostreatus</i> , <i>Trametes versicolour</i> (formerly known as <i>Coriolus versicolor</i> or <i>Polyporus versicolor</i>) <i>Trametes versicolor</i> <i>Tramete. pubescens</i> , <i>T. hirsuta</i> <i>T. gallica</i> <i>Trichoderma atroviride</i> , <i>T. harzianum</i> , <i>T. longibrachiatum</i> <i>Pleurotus ostreatus</i> , <i>Pycnoporus sanguineus</i>	<ul style="list-style-type: none"> • delignification of lignocellulosic material, • sporulation, • pigment production, • fungal morphogenesis, • protection against toxic compounds, • fruiting body formation • plant pathogenesis 	[1,13,14,15,16,17,18,7,19,20]
Bacteria	<i>Azospirillum lipoferum</i> <i>Azospirillum lipoferum</i> , <i>Bacillus subtilis</i> , <i>Streptomyces lavendulae</i> , <i>S. cyaneus</i> <i>Marinomonas mediterranea</i> . <i>Bacillus subtilis</i> <i>Streptomyces cyaneus</i> ,	<ul style="list-style-type: none"> • melanin production for cell pigmentation • spore coat resistance against hydrogen peroxide and UV • involvement in morphogenesis • involved in the solubilization and mineralization of lignin 	[21,18,14,22]

Table 2: Production of laccase under solid state fermentation

Source	Support	Reactor	Production (U/l)	Reference
<i>Galerina</i> sp. HCl	Orange peels	Cotton plugged Erlenmeyer flasks	720	Laura Mendoza et al. 2010
<i>Pycnoporus cinnabarinus</i>	Sugarcane bagasse	Packed reactor	10000	Meza et al. 2006
<i>Trametes hirsuta</i>	Grape seeds	Tray	18715	Rodríguez Couto et al. 2006
<i>Trametes hirsuta</i>	Kiwi fruits	Cotton plugged Erlenmeyer flasks	5399	Rosales et al. 2005
<i>Trametes hirsuta</i>	Nylon sponge	Tray	6898	Rodríguez Couto et al. 2006
<i>Trametes hirsuta</i>	Orange peels	Tray	12000	Rosales et al. 2007
<i>Trametes pubescens</i>	Banana skin	Cotton plugged Erlenmeyer flasks	1570	Osma et al. 2007
<i>Trametes versicolor</i>	Barley bran	Immersion	600	Rodríguez Couto et al. 2003
<i>Trametes versicolor</i>	Barley bran	Tray	3500	Rodríguez Couto et al. 2003
<i>Trametes versicolor</i>	Nylon sponge	Expanded bed	229	Rodríguez Couto et al. 2003
<i>Trametes versicolor</i>	Nylon sponge	Immersion	229	Rodríguez Couto et al. 2003
<i>Trametes versicolor</i>	Nylon sponge	Tray	343	Rodríguez Couto et al. 2003

Table 3 : Commercial preparations based on laccases for industrial processes.

Industry	Main application	Brand name	Manufacturer
Food industry	Brewing	Flavourstar	Advanced Enzyme Technologies Ltd. (India)
	Colour enhancement in tea, etc.	LACCASE Y120	Amano Enzyme USA Co. Ltd.
	Cork modification	Suberase	Novozymes (Denmark)
Paper industry	Pulp bleaching	Lignozym-process	Lignozym GmbH (Germany)
	Paper pulp delignification	Novozym 51003	Novozymes (Denmark)
Textile Industry	Denim bleaching	Bleach Cut 3-S	Season Chemicals (China)
		DeniLite	Novozymes (Denmark)
		ZyLite	Zytext Pvt. Ltd. (India)
	Denim finishing	Cololacc BB	Colotex Biotechnology Co. Ltd. (Hong Kong)
		Ecoston LC10	AB Enzymes GmbH (Germany)
		IndiStar	Genencor Inc. (Rochester, USA)
		Novoprime Base 268	Novozymes (Denmark)
Denim bleaching and shading	Primagreen Ecofade LT100	Genencor Inc. (Rochester, USA)	

5. Conclusion

Laccases are widespread in nature, being produced by a wide variety of plants, fungi and also bacteria. The functions of the enzyme differ from organism to organism to organisms. Laccases catalyse the oxidation of phenolic compounds whilst simultaneously reducing molecular oxygen to water. The catalytic ability of laccases has, not surprisingly, led to diverse biotechnological applications of this enzyme. The introduction of the laccase-mediator system provides a biological alternative to traditional bleaching processes. The laccase enzyme has a wide field of application including food industries, the pulp and paper industries, the treatment of various industrial effluents, enzymatic decolouring of material and bioremediation of soils. It is therefore not surprising that this enzyme has

been studied intensively since the nineteenth century and yet remains a topic of intense research today.

6. Reference

1. Sudhir Kumar and Abha Mishra, (2011) Optimization of laccase production from WRF-1 on groundnut shell and cyanobacterial biomass: By application of Box-Behnken experimental design, J. Microbiol. Biotech. Res., 2011, 1 (2): 33-53.
2. Y.Arakane, S.Muthukrishnan, R.W.Beeman, M.R.Kanost, K.J.Kramer, (2005). "Laccase 2 is the phenoloxidase gene required for beetle cuticle tanning," Proceedings of The National Academy

- of Sciences of the United States of America, vol. 102, no. 32, pp. 11337–11342.
3. Alpeshkumar J. Shiroya, recent advances of laccase enzyme in industrial biotechnology: a review, *pharmatutor-art-1419* *pharmatutor-art-1419*
 4. A.E. Gardiol, R.J. Hernandez, B.R. Harte, (1998). Device for detecting oxygen with oxidase. United States Patent 5,804,401.
 5. Y. Arakane, S. Muthukrishnan, R.W. Beeman, M.R. Kanost, K.J. Kramer, (2005). “Laccase 2 is the phenoloxidase gene required for beetle cuticle tanning,” *Proceedings of The National Academy of Sciences of the United States of America*, vol. 102, no. 32, pp. 11337–11342.
 6. A. De Marco, K.A. Roubelakis-Angelakis, (1997). Laccase activity could contribute to cell-wall reconstitution in regenerating protoplasts. *Phytochemistry*, 46:421.425.
 7. D.M. O’Malley, R. Whetten, W. Bao, C.L. Chen, R.R. Sederoff, (1993). The role of laccase in lignification. *Plant J.*, 4:751.757.
 8. E. Lehman, E. Harel, A.M. Mayer, (1974). Copper content and other characteristics of purified peach laccase. *Phytochemistry*, 13:1713.1717.
 9. P. Ranocha, G. McDougall, S. Hawkins, R. Sterjiades, G. Borderies, D. Stewart, M. Cabanes-Macheteau, A.M. Boudet, D. Goffner, (1999). Biochemical characterization, molecular cloning and expression of laccases a divergent gene family - in poplar. *Eur. J. Biochem*, 259:485.495.
 10. R. Bligny, R. Douce, (1983). Excretion of laccase by sycamore (*Acer pseudoplatanus* L.) cells. Purification and properties of the enzyme. *Biochem. J.*, 209:489.496.
 11. U. Dwivedi, P. Singh, V. Pandey, A. Kumar, (2011). *J Mol Catal B- Enzym*, 68: 117-128.
 12. W.G. Levine, (1965). Laccase: a review, In: *The biochemistry of copper* (Academic Press Inc., New York), pp. 371-385.
 13. C. Galhaup, H. Wagner, B. Hinterstoisser, D. Haltrich, (2002). Increased production of laccase by the wood-degrading basidiomycete *Trametes pubescens*. *Enz. Microb. Technol.*, 30:529.536.
 14. C. Held, A. Kandelbauer, M. Schroeder, A. Cavaco-Paulo G. Guebitz, (2005). *Environ Chem Lett*, 3: 74–77.
 15. C.S. Evans and J.N. Hedger, Degradation of plant cell wall polymers, *Fungi in bioremediation*. (Ed. G.M. Gadd, British Mycological Society. Cambridge Univ. Press. UK 2001), pp. 1-20.
 16. C.S. Chen, M. Mrkisch, S. Huang, G.M. Whitesides, D.E. Ingber, (1998). Micropatterned surfaces for control of cell shape, position, and function. *Biotechnol Prog*, 14:356–63.
 17. D.D’Souza-Ticlo, D. Sharma, C. Raghukumar, (2009). “A thermostable metal-tolerant laccase with bioremediation potential from a marine-derived fungus,” *Marine Biotechnology*, vol. 11, no. 6, pp. 725–737.
 18. D. Faure, M.L. Bouillant, R. Bally, (1994). Isolation of *Azospirillum lipoferum* 4T Tn5 mutants affected in melanization and laccase activity. *Appl. Environ. Microbiol.*, 60:3413.3415.
 19. D. Rigling, N.K. van Alfen, (1991). Regulation of laccase biosynthesis in the plant-pathogenic fungus *Cryphonectria parasitica* by double-stranded RNA. *J. Bacteriol*, 173:8000.8003.
 20. D.S. Yaver, R.M. Berka, S.H. Brown, F. Xu, (2001). The presymposium on recent advances in lignin biodegradation and biosynthesis, Vikki Biocentre, University of Helsinki, Finland, pp.40.
 21. A. Givaudan, A. Effosse, D. Faure, P. Potier, M.L. Bouillant, R. Bally, (1993). Polyphenol oxidase in *Azospirillum lipoferum* isolated from rice rhizosphere: evidence for laccase activity in non-motile strains of *Azospirillum lipoferum*. *FEMS Microbiol. Lett.*, 108:205.210.
 22. P. Sharma, R. Goel N. Capalash, (2007). *World J Microbiol Biotechnol*, 23:823–832.
 23. A. Kunamneni, F. Plou, A. Ballesteros, M. Alcalde, (2008). *Recent Patents on Biotechnology* 2: 10-24.
 24. F.S. Archibald, R. Bourbonnais, L. Jurasek, M.G. Paice, and I.D. Reid, *Journal of Biotechnology*, 53, 215 (1997).
 25. M. Heinzkill, L. Bech, T. Halkier, P. Schneider and T. Anke, *Applied and Environmental Microbiology* 64, 1601 (1998).
 26. A.M.R.B. Xavier, D.V. Evtuguin, R.M.P. Ferreira and F.L. Amado, Laccase production for lignin oxidative activity. *Proceedings of the 8th International Conference on Biotechnology in the Pulp and Paper Industry*, 4-8 June, Helsinki, Finland, 2001.
 27. J.A. Buswell, Y. Cai and S. Chang, *FEMS Microbiology Letters* 128, 81 (1995). H. Park, *Biotechnology Letters* 21, 965 (1999).
 28. V. Faraco, P. Giardina, G. Palmieri and G. Sannia, *Progress in Biotechnology* 21, 105 (2002).
 29. G. Songulashvili, V. Elisashvili, S. Wasser, E. Nevo, Y. Hadar, (2006). *Biotechnol Lett*, 28:1425–1429.
 30. Robene-Soustrade and B. Lung-Escarmant, *European Journal of Forest Pathology* 27, 105 (1997).
 31. N. Hatvani, I. Mécs, (2001). *Process Biochem*, 37: 491-496.
 32. M. Staji, L. Persky, D. Friesem, Y. Hadar, S. Wasser, E. Nevo, et al., (2006). *Enzyme Microb Technol*, 38: 65-73.
 33. M.C. Monteiro and M.E.A. De Carvalho, *Applied Biochemistry and Biotechnology* 70-72, 983 (1998).
 34. C. Galhaup and D. Haltrich, *Applied Microbiology and Biotechnology* 56, 225 (2001).
 35. A.F. Vasconcelos, A.M. Barbosa, R.F.H. Dekker, I.S. Scarminio and M.I. Rezende, *Process Biochemistry* 35, 1131 (2000).
 36. E. Selinheimo, K. Kruus, J. Buchert, A. Hopia, K. Autio, (2006). Effects of laccase, xylanase and

- their combination on the rheological properties of wheat doughs. *J Cereal Sci*, 43:152–9.
37. S.Khammuang, R.Sarnthima, (2009). “Laccase activity from fresh fruiting bodies of *Ganoderma* sp. MK05: purification and remazol brilliant blue R decolorization,” *Journal of Biological Sciences*, vol. 9, no. 1, pp. 83–87.
 38. R.F.Taylor, (1991). *Protein immobilization: Fundamental and applications* (Marcel Dekker Inc., New York, USA).
 39. R.C.Minussi, G.M.Pastore, N.Durán, (2002). Potential applications of laccase in the food industry. *Trends Food Sci Technol*, 13:205–16.
 40. A.I. Yaropolov, O.V. Skorobogat'ko, S.S. Vartanov and S.D. Varfolomeyev, *Applied Biochemistry and Biotechnology* 49, 257 (1994).
 41. P.T.Hammond, G.M.Whitesides, (1995). Formation of polymer microstructures by selective deposition of polyion multilayers using patterned selfassembled monolayers as a template. *Macromolecules*, 28: 7569–71.